



Figure 7.2 Cleavage of a frog egg



DEVELOPMENTAL BIOLOGY, 9e, Figure 7.2

© 2010 Sinauer Associates, Inc.


Figure 7.3 Scanning electron micrographs of frog egg cleavage



DEVELOPMENTAL BIOLOGY, 9e, Figure 7.3

© 2010 Sinauer Associates, Inc.


Figure 7.5 Fate maps of the *Xenopus laevis* blastula exterior (A) and interior (B)



DEVELOPMENTAL BIOLOGY, 9e, Figure 7.5

© 2010 Sinauer Associates, Inc.


Figure 7.6 Cell movements during frog gastrulation (Part 1)



DEVELOPMENTAL BIOLOGY, 9e, Figure 7.6 (Part 1)

© 2010 Sinauer Associates, Inc.


Figure 7.6 Cell movements during frog gastrulation (Part 2)



DEVELOPMENTAL BIOLOGY, 9e, Figure 7.6 (Part 2)

© 2010 Sinauer Associates, Inc.


Figure 7.6 Cell movements during frog gastrulation (Part 3)



DEVELOPMENTAL BIOLOGY, 9e, Figure 7.6 (Part 3)

© 2010 Sinauer Associates, Inc.


Figure 7.6 Cell movements during frog gastrulation (Part 4)



DEVELOPMENTAL BIOLOGY, 9e, Figure 7.6 (Part 4)

© 2010 Sinauer Associates, Inc.

Figure 7.7 Surface view of an early dorsal blastopore lip of *Xenopus*



DEVELOPMENTAL BIOLOGY, 9e, Figure 7.7

© 2010 Sinauer Associates, Inc.



Figure 7.11 Protocadherin expression separates axial and paraxial mesoderm




Figure 12.1 Mesodermal development in frog and chick embryos






Figure 8.15 Development of a human embryo from fertilization to implantation



DEVELOPMENTAL BIOLOGY, 9e, Figure 8.15

© 2010 Sinauer Associates, Inc.


Figure 8.23 Amnion structure and cell movements during human gastrulation




DEVELOPMENTAL BIOLOGY, 9e, Figure 8.23

© 2010 Sinauer Associates, Inc.



Figure 12.1 Mesodermal development in frog and chick embryos





Primitive streak stage (7.5 dpc)







Chapman DL, Aguirre J, Hancock S, Shivel LM, Papaioannou VL. *Tbx6*, a mouse *T-box* gene implicated in paraxial mesoderm formation at gastrulation. *Dev Biol.* 1996 Dec 15;180(2):534-42.













Summary



Chapman DL, Papaioannou VE. Three neural tubes in mouse embryos with mutations in the T-box gene Tbx6. *Nature*. 1998 Feb 12;391(6668):695-7.

Figure 7.6 Cell movements during frog gastrulation (Part 3)



DEVELOPMENTAL BIOLOGY, 9e, Figure 7.6 (Part 3)

© 2010 Sinauer Associates, Inc.