



6.16 The sonic hedgehog gene is shown by in situ hybridization to be expressed in the chick nervous system (red arrow), gut (blue arrow), and limb bud (black arrow) of a 3-day chick embryo







DEVELOPMENTAL BIOLOGY, Eighth Edition, Figure 6.16 © 2006 Sinauer Associates, Inc.



4.18 Production of transgenic mice (Part 1)




DEVELOPMENTAL BIOLOGY, Eighth Edition, Figure 4.18 (Part 1) © 2006 Sinauer Associates, Inc.







4.20 Morphological analysis of *Bmp7* knockout mice (Part 1)



DEVELOPMENTAL BIOLOGY, Eighth Edition, Figure 4.20 (Part 1) © 2006 Sinauer Associates, Inc.

ARTICLES

**Cyclopia and defective axial patterning in mice lacking *Sonic hedgehog* gene function**

Chin Chiang\*†, Ying Litingtung\*†, Eric Lee\*, Keith E. Young†, Jeffrey L. Corden†, Heiner Westphal\* & Philip A. Beachy†

\* Laboratory of Mammalian Genes and Development, National Institute of Health, Bethesda, Maryland 20892, USA

† Howard Hughes Medical Institute, Department of Molecular Biology and Genetics, Johns Hopkins University School of Medicine, Baltimore, Maryland 21205, USA

